



PATENT APPLICATION TRANSMITTAL

(Only for nonprovisional applications under 37 CFR § 1.53(b))

Attorney Docket No.

920010.448C8

First Inventor or Application Identifier

Martin A. Cheever

Title

COMPOUNDS FOR ELICITING OR ENHANCING IMMUNE REACTIVITY
TO HER-2/neu PROTEIN FOR PREVENTION OR TREATMENT OF
MALIGNANCIES WHICH THE HER-2/neu ONCOGENE IS ASSOCIATED

Express Mail Label No.

EM001237621US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

1. ☐ General Authorization Form & Fee Transmittal
(Submit an original and a duplicate for fee processing)

2. ☒ Specification [Total Pages] **66**
(preferred arrangement set forth below)
- Descriptive Title of the Invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings (if filed)
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure

3. ☒ Drawing(s) (35 USC 113) [Total Sheets] **4**

4. ☐ Oath or Declaration [Total Pages] **2**

- a. ☐ Newly executed (original or copy)
- b. ☒ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
- i. ☐ **DELETION OF INVENTOR(S)**
Signed statement attached deleting
inventor(s) named in the prior
application,
see 37 CFR 1.63(d)(2) and 1.33(b)

5. ☒ Incorporation By Reference (useable if box 4b is checked)
The entire disclosure of the prior application, from which
a copy of the oath or declaration is supplied under Box
4b, is considered to be part of the disclosure of the
accompanying application and is hereby incorporated by
reference therein.

6. ☐ Microfiche Computer Program (Appendix)

7. Nucleotide and Amino Acid Sequence Submission
(if applicable, all necessary)

- a. ☐ Computer-Readable Copy
- b. ☐ Paper Copy (identical to computer copy)
- c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement (when there is an assignee) ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard
14. ☐ Small Entity Statement(s) ☒ Statement filed in prior application, Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. ☒ Other: Certificate of Express Mail

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information below and in a preliminary amendment

☒ Continuation ☐ Divisional ☐ Continuation-In-Part (CIP) of prior Application No.: 08/625,101

Prior application information: Examiner N. JohnsonGroup / Art Unit 1642

☐ Claims the benefit of Provisional Application No. _____

CORRESPONDENCE ADDRESS

Richard G. Sharkey
Seed and Berry LLP
6300 Columbia Center
701 Fifth Avenue
Seattle, Washington 98104-7092
(206) 622-4900 phone
(206) 682-6031 fax

Respectfully submitted,

TYPED OR PRINTED NAME Richard G. SharkeyREGISTRATION NO. 32,629SIGNATURE Richard G. SharkeyDate October 6, 1998

u:\float\gac\0307

Description

COMPOUNDS FOR ELICITING OR ENHANCING IMMUNE REACTIVITY
TO HER-2/*neu* PROTEIN FOR PREVENTION OR TREATMENT OF
5 MALIGNANCIES IN WHICH THE HER-2/*neu* ONCOGENE IS ASSOCIATED

Cross Reference to Related Application

This application is a continuation-in-part to
Serial No. 414,417, filed March 31, 1995, which is a
10 continuation-in-part application to Serial No. 106,112,
filed August 12, 1993, abandoned, which is a continuation-
in-part application to Serial No. 033,644, filed March 17,
1993, abandoned.

15 Technical Field

The present invention is generally directed
toward polypeptides, and nucleic acid molecules encoding
such polypeptides, for eliciting or enhancing an immune
response to HER-2/*neu* protein, including for use in the
20 treatment of malignancies in which the HER-2/*neu* oncogene
is associated.

Background of the Invention

Despite enormous investments of financial and
25 human resources, cancer remains one of the major causes of
death. For example, cancer is the leading cause of death
in women between the ages of 35 and 74. Breast cancer is
the most common malignancy in women and the incidence for
developing breast cancer is on the rise. One in nine
30 women will be diagnosed with the disease. Standard
approaches to cure breast cancer have centered around a
combination of surgery, radiation and chemotherapy. These
approaches have resulted in some dramatic successes in
certain malignancies. However, these approaches have not

09157316-100500

been successful for all malignancies and breast cancer is most often incurable when attempting to treat beyond a certain stage. Alternative approaches to prevention and therapy are necessary.

5 A common characteristic of malignancies is uncontrolled cell growth. Cancer cells appear to have undergone a process of transformation from the normal phenotype to a malignant phenotype capable of autonomous growth. Amplification and overexpression of somatic cell
10 genes is considered to be a common primary event that results in the transformation of normal cells to malignant cells. The malignant phenotypic characteristics encoded by the oncogenic genes are passed on during cell division to the progeny of the transformed cells.

15 Ongoing research involving oncogenes has identified at least forty oncogenes operative in malignant cells and responsible for, or associated with, transformation. Oncogenes have been classified into different groups based on the putative function or
20 location of their gene products (such as the protein expressed by the oncogene).

 Oncogenes are believed to be essential for certain aspects of normal cellular physiology. In this regard, the HER-2/*neu* oncogene is a member of the tyrosine
25 protein kinase family of oncogenes and shares a high degree of homology with the epidermal growth factor receptor. HER-2/*neu* presumably plays a role in cell growth and/or differentiation. HER-2/*neu* appears to induce malignancies through quantitative mechanisms that
30 result from increased or deregulated expression of an essentially normal gene product.

 HER-2/*neu* (p185) is the protein product of the HER-2/*neu* oncogene. The HER-2/*neu* gene is amplified and the HER-2/*neu* protein is overexpressed in a variety of

09167516-100698

cancers including breast, ovarian, colon, lung and prostate cancer. HER-2/*neu* is related to malignant transformation. It is found in 50%-60% of ductal *in situ* carcinoma and 20%-40% of all breast cancers, as well as a substantial fraction of adenocarcinomas arising in the ovaries, prostate, colon and lung. HER-2/*neu* is intimately associated not only with the malignant phenotype, but also with the aggressiveness of the malignancy, being found in one-fourth of all invasive breast cancers. HER-2/*neu* overexpression is correlated with a poor prognosis in both breast and ovarian cancer. HER-2/*neu* is a transmembrane protein with a relative molecular mass of 185 kd that is approximately 1255 amino acids (aa) in length. It has an extracellular binding domain (ECD) of approximately 645 aa, with 40% homology to epidermal growth factor receptor (EGFR), a highly hydrophobic transmembrane anchor domain (TMD), and a carboxyterminal cytoplasmic domain (CD) of approximately 580 aa with 80% homology to EGFR.

Due to the difficulties in the current approaches to therapy of cancers in which the HER-2/*neu* oncogene is associated, there is a need in the art for improved compounds and compositions. The present invention fulfills this need, and further provides other related advantages.

Summary of the Invention

Briefly stated, the present invention provides polypeptides, nucleic acid molecules (directing the expression of such polypeptides) and viral vectors (directing the expression of such polypeptides) for uses which include the immunization of a warm-blooded animal against a malignancy in which the HER-2/*neu* oncogene is associated. A polypeptide or nucleic acid molecule

00167516-100608

according to this invention may be present in a composition that includes a pharmaceutically acceptable carrier or diluent. Such a polypeptide, nucleic acid molecule, viral vector or pharmaceutical composition may
5 be administered on a one-time basis (e.g., when a malignancy is suspected) or on a periodic basis (e.g., for an individual with an elevated risk of acquiring or reacquiring a malignancy). A compound or composition of the present invention may be useful in the treatment of an
10 existing tumor or to prevent tumor occurrence or reoccurrence.

In one aspect, the present invention provides compounds and compositions that elicit or enhance an immune response to HER-2/*neu* protein. One embodiment of
15 the present invention provides a polypeptide encoded by a DNA sequence selected from: (a) nucleotides 2026 through 3765 of SEQ ID NO:1; and (b) DNA sequences that hybridize to a nucleotide sequence complementary to nucleotides 2026 through 3765 of SEQ ID NO:1 under moderately stringent
20 conditions, wherein the DNA sequence encodes a polypeptide that produces an immune response to HER-2/*neu* protein. In a preferred embodiment, a polypeptide has the amino acid sequence of SEQ ID NO:2 from lysine, amino acid 676, through valine, amino acid 1255, or a variant thereof that
25 produces at least an equivalent immune response. A composition is provided that comprises a polypeptide of the present invention in combination with a pharmaceutically acceptable carrier or diluent. In another embodiment, a nucleic acid molecule directing the
30 expression of a polypeptide according to the present invention is provided. In another embodiment, a viral vector directing the expression of a polypeptide according to the present invention is provided.

09167516.100698

09167516.100698

In another aspect, the present invention provides a method for eliciting or enhancing an immune response to HER-2/*neu* protein, comprising administering to a warm-blooded animal (such as a human) in an amount effective to elicit or enhance the response a polypeptide according to the present invention, or a nucleic acid molecule or a viral vector, either directing the expression of such a polypeptide. In one embodiment, a peptide is administered in combination with a pharmaceutically acceptable carrier or diluent. In another embodiment, the step of administering comprises transfecting cells of the animal *ex vivo* with the nucleic acid molecule and subsequently delivering the transfected cells to the animal. In another embodiment, the step of administering comprises infecting cells of the animal *ex vivo* with the viral vector and subsequently delivering the infected cells to the animal.

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings.

Brief Description of the Drawings

Figure 1 shows the results of the priming of naive T lymphocytes to HER-2/*neu* polypeptide by dendritic cells. Bone marrow-derived DC were generated with GM-CSF and IL6 from CD34+ stem cells. DC pulsed with HER-2/*neu* polypeptide induced protein-specific proliferation of autologous CD4+/CD45RA+ T lymphocytes after 7 days of culturing T cells with DC. Bone marrow-derived CD34+ stem cells cultured for one week in serum-free medium containing GM-CSF and IL-6 were used as APC. APC were plated into 96-well round-bottomed plates (Corning, Corning, NY, USA) at various concentrations and incubated for 16-18 hours with 20-25 µg/ml of recombinant HER-2/*neu*

polypeptide. CD4+ T lymphocytes were isolated from autologous peripheral blood mononuclear cells by positive selection using immunoaffinity columns (CellPro, Inc., Bothell, WA, USA). Antigen-pulsed APC were irradiated (10 Gy), and CD4+ T lymphocytes were added at 10^5 per well. Proliferative response of T cells was measured by the uptake of (^3H)thymidine ($1\mu\text{Ci}/\text{well}$) added on day 7 for 16-18 hours. Proliferation assays were performed in serum- and cytokine-free medium in 5 well replicates. The symbols represent: \bullet DC + HER-2/*neu* polypeptide + CD4+/CD45RA+ T cells; \circ DC + CD4+/CD45RA+ T cells; and \square DC + HER-2/*neu* polypeptide.

Figure 2 shows the response of CD4+ cells to HER-2/*neu* polypeptide. Using the priming assay described for Figure 1, CD4+ T cells from normal donors were tested for responses to recombinant human HER-2/*neu* polypeptide. The symbols represent: \blacksquare SC+CD4; and \blacklozenge SC+CD4+HER-2/*neu* polypeptide. "SC" is stem cells.

Figure 3 shows that rats immunized with rat HER-2/*neu* polypeptide develop rat *neu* specific antibodies. Rats were immunized with recombinant rat HER-2/*neu* polypeptide 25 ug in MPL or vaxcel adjuvant. Three immunizations were given, each 20 days apart. Twenty days after the final immunization rats were assessed for antibody responses to rat *neu*. Animals immunized with rat HER-2/*neu* polypeptide and the vaxcel adjuvant showed high titer rat *neu* specific responses. The control was an animal immunized with human HER-2/*neu* polypeptide (foreign protein). In separate experiments, rats immunized with 100 ug and 300 ug of purified whole rat *neu* did not develop detectable *neu* specific antibodies (data not shown). Data represents the mean and standard deviation of 3 animals. The symbols represent: \blacksquare rat HER-2/*neu*

09167516.100698

polypeptide/MPL; ● rat HER-2/neu polypeptide/vaccel;
 □ MPL alone; ○ vaccel alone; and Φ control.
 "MPL" and "vaccel" are adjuvants (Ribi, Bozeman, MT, USA).
 "Neu" is HER-2/neu protein.

- 5 Figure 4 shows that breast cancer patients have
 preexistent immunity to HER-2/neu polypeptide. Patient
 PBMC were evaluated by tritiated thymidine incorporation
 in 24 well replicates. Responsive wells are scored as
 greater than the mean and 3 standard deviations (372 cpm)
 10 of the control wells. This HER-2/neu positive-stage II
 breast cancer patient has a significant response to
 recombinant human HER-2/neu polypeptide. The symbols "p"
 represent peptides for HER-2/neu protein, "tt" represents
 tetanus toxoid, and "hHNP" represents recombinant human
 15 HER-2/neu polypeptide.

Detailed Description of the Invention

- Prior to setting forth the invention, it may be
 helpful to an understanding thereof to set forth
 20 definitions of certain terms to be used hereinafter.

- HER-2/neu polypeptide - as used herein, refers
 to a portion of the HER-2/neu protein (the protein also
 known as p185 or c-erbB2) having the amino acid sequence
 of SEQ ID NO:2 from lysine, amino acid 676, through
 25 valine, amino acid 1255; and may be naturally derived,
 synthetically produced, genetically engineered, or a
 functionally equivalent variant thereof, e.g., where one
 or more amino acids are replaced by other amino acid(s) or
 non-amino acid(s) which do not substantially affect
 30 elicitation or enhancement of an immune response to HER-
 2/neu protein (e.g., variant stimulates a response by
 helper T cells or cytotoxic T cells).

Proliferation of T cells - as used herein,
 includes the multiplication of T cells as well as the

00167516-100658

stimulation of T cells leading to multiplication, i.e., the initiation of events leading to mitosis and mitosis itself. Methods for detecting proliferation of T cells are discussed below.

5 As noted above, the present invention is directed toward compounds and compositions to elicit or enhance immunity to the protein product expressed by the HER-2/neu oncogene, including for malignancies in a warm-blooded animal wherein an amplified HER-2/neu gene is
10 associated with the malignancies. Association of an amplified HER-2/neu gene with a malignancy does not require that the protein expression product of the gene be present on the tumor. For example, overexpression of the protein expression product may be involved with initiation
15 of a tumor, but the protein expression may subsequently be lost. A use of the present invention is to elicit or enhance an effective autochthonous immune response to convert a HER-2/neu positive tumor to HER-2/neu negative.

More specifically, the disclosure of the present
20 invention, in one aspect, shows that a polypeptide based on a particular portion (HER-2/neu polypeptide) of the protein expression product of the HER-2/neu gene can be recognized by thymus-dependent lymphocytes (hereinafter "T cells") and, therefore, the autochthonous immune T cell
25 response can be utilized prophylactically or to treat malignancies in which such a protein is or has been overexpressed. The disclosure of the present invention also shows, in another aspect, that nucleic acid molecules directing the expression of such a peptide may be used
30 alone or in a viral vector for immunization.

In general, CD4⁺ T cell populations are considered to function as helpers/inducers through the release of lymphokines when stimulated by a specific antigen; however, a subset of CD4⁺ cells can act as

09167516-100698

cytotoxic T lymphocytes (CTL). Similarly, CD8⁺ T cells are considered to function by directly lysing antigenic targets; however, under a variety of circumstances they can secrete lymphokines to provide helper or DTH function.

- 5 Despite the potential of overlapping function, the phenotypic CD4 and CD8 markers are linked to the recognition of peptides bound to class II or class I MHC antigens. The recognition of antigen in the context of class II or class I MHC mandates that CD4⁺ and CD8⁺ T
10 cells respond to different antigens or the same antigen presented under different circumstances. The binding of immunogenic peptides to class II MHC antigens most commonly occurs for antigens ingested by antigen presenting cells. Therefore, CD4⁺ T cells generally
15 recognize antigens that have been external to the tumor cells. By contrast, under normal circumstances, binding of peptides to class I MHC occurs only for proteins present in the cytosol and synthesized by the target itself, proteins in the external environment are excluded.
20 An exception to this is the binding of exogenous peptides with a precise class I binding motif which are present outside the cell in high concentration. Thus, CD4⁺ and CD8⁺ T cells have broadly different functions and tend to recognize different antigens as a reflection of where the
25 antigens normally reside.

- As disclosed within the present invention, a polypeptide portion of the protein product expressed by the HER-2/*neu* oncogene is recognized by T cells. Circulating HER-2/*neu* polypeptide is degraded to peptide
30 fragments. Peptide fragments from the polypeptide bind to major histocompatibility complex (MHC) antigens. By display of a peptide bound to MHC antigen on the cell surface and recognition by host T cells of the combination of peptide plus self MHC antigen, HER-2/*neu* polypeptide

00167515.100698

(including that expressed on a malignant cell) will be immunogenic to T cells. The exquisite specificity of the T cell receptor enables individual T cells to discriminate between peptides which differ by a single amino acid residue.

During the immune response to a peptide fragment from the polypeptide, T cells expressing a T cell receptor with high affinity binding of the peptide-MHC complex will bind to the peptide-MHC complex and thereby become activated and induced to proliferate. In the first encounter with a peptide, small numbers of immune T cells will secrete lymphokines, proliferate and differentiate into effector and memory T cells. The primary immune response will occur *in vivo* but has been difficult to detect *in vitro*. Subsequent encounter with the same antigen by the memory T cell will lead to a faster and more intense immune response. The secondary response will occur either *in vivo* or *in vitro*. The *in vitro* response is easily gauged by measuring the degree of proliferation, the degree of cytokine production, or the generation of cytolytic activity of the T cell population re-exposed in the antigen. Substantial proliferation of the T cell population in response to a particular antigen is considered to be indicative of prior exposure or priming to the antigen.

The compounds of this invention generally comprise HER-2/*neu* polypeptides or DNA molecules that direct the expression of such peptides, wherein the DNA molecules may be present in a viral vector. As noted above, the polypeptides of the present invention include variants of the polypeptide of SEQ ID NO:2 from amino acid 676 through amino acid 1255, that retain the ability to stimulate an immune response. Such variants include various structural forms of the native polypeptide. Due

00157515-100598

to the presence of ionizable amino and carboxyl groups, for example, a HER-2/*neu* polypeptide may be in the form of an acidic or basic salt, or may be in neutral form. Individual amino acid residues may also be modified by
5 oxidation or reduction.

Variants within the scope of this invention also include polypeptides in which the primary amino acid structure native HER-2/*neu* polypeptide is modified by forming covalent or aggregative conjugates with other
10 peptides or polypeptides, or chemical moieties such as glycosyl groups, lipids, phosphate, acetyl groups and the like. Covalent derivatives may be prepared, for example, by linking particular functional groups to amino acid side chains or at the N- or C-terminus.

The present invention also includes HER-2/*neu* polypeptides with or without glycosylation. Polypeptides expressed in yeast or mammalian expression systems may be similar to or slightly different in molecular weight and glycosylation pattern than the native molecules, depending
15 upon the expression system. For instance, expression of DNA encoding polypeptides in bacteria such as *E. coli* typically provides non-glycosylated molecules. N-glycosylation sites of eukaryotic proteins are characterized by the amino acid triplet Asn-A₁-Z, where A₁
20 is any amino acid except Pro, and Z is Ser or Thr. Variants of HER-2/*neu* polypeptides having inactivated N-glycosylation sites can be produced by techniques known to those of ordinary skill in the art, such as oligonucleotide synthesis and ligation or site-specific
25 mutagenesis techniques, and are within the scope of this invention. Alternatively, N-linked glycosylation sites can be added to a HER-2/*neu* polypeptide.
30

The polypeptides of this invention also include variants of the SEQ ID NO:2 polypeptide (i.e., variants of

00167516-100698

a polypeptide having the amino acid sequence of SEQ ID NO:2 from amino acid 676 through amino acid 1255) that have an amino acid sequence different from this sequence because of one or more deletions, insertions, 5 substitutions or other modifications. In one embodiment, such variants are substantially homologous to the native HER-2/*neu* polypeptide and retain the ability to stimulate an immune response. "Substantial homology," as used herein, refers to amino acid sequences that may be encoded 10 by DNA sequences that are capable of hybridizing under moderately stringent conditions to a nucleotide sequence complimentary to a naturally occurring DNA sequence encoding the specified polypeptide portion of SEQ ID NO:2 herein (i.e., nucleotides 2026 through 3765 of SEQ ID 15 NO:1). Suitable moderately stringent conditions include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC (containing 0.1% SDS). Such 20 hybridizing DNA sequences are also within the scope of this invention. The effect of any such modifications on the ability of a HER-2/*neu* polypeptide to produce an immune response may be readily determined (e.g., by analyzing the ability of the mutated HER-2/*neu* polypeptide 25 to induce a T cell response using, for example, the methods described herein).

Generally, amino acid substitutions may be made in a variety of ways to provide other embodiments of variants within the present invention. First, for 30 example, amino acid substitutions may be made conservatively; i.e., a substitute amino acid replaces an amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the

polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; 5 (4) lys, arg, his; and (5) phe, tyr, trp, his. An example of a non-conservative change is to replace an amino acid of one group with an amino acid from another group.

Another way to make amino acid substitutions to produce variants of the present invention is to identify 10 and replace amino acids in T cell motifs with potential to bind to class II MHC molecules (for CD4+ T cell response) or class I MHC molecules (for CD8+ T cell response). Peptide segments (of a HER-2/neu polypeptide) with a motif with theoretical potential to bind to class II MHC 15 molecules may be identified by computer analysis. For example, a protein sequence analysis package, *T Sites*, that incorporates several computer algorithms designed to distinguish potential sites for T cell recognition can be used (Feller and de la Cruz, *Nature* 349:720-721, 1991). 20 Two searching algorithms are used: (1) the AMPHI algorithm described by Margalit (Feller and de la Cruz, *Nature* 349:720-721, 1991; Margalit et al., *J. Immunol.* 138:2213-2229, 1987) identifies epitope motifs according to alpha-helical periodicity and amphipathicity; (2) the 25 Rothbard and Taylor algorithm identifies epitope motifs according to charge and polarity pattern (Rothbard and Taylor, *EMBO* 7:93-100, 1988). Segments with both motifs are most appropriate for binding to class II MHC molecules. CD8+ T cells recognize peptide bound to class 30 I MHC molecules. Falk et al. have determined that peptides binding to particular MHC molecules share discernible sequence motifs (Falk et al., *Nature* 351:290-296, 1991). A peptide motif for binding in the groove of HLA-A2.1 has been defined by Edman degradation of peptides

stripped from HLA-A2.1 molecules of a cultured cell line (Table 2, from Falk et al., *supra*). The method identified the typical or average HLA-A2.1 binding peptide as being 9 amino acids in length with dominant anchor residues occurring at positions 2 (L) and 9 (V). Commonly occurring strong binding residues have been identified at positions 2 (M), 4 (E,K), 6 (V), and 8 (K). The identified motif represents the average of many binding peptides.

The HLA-A2.1 Restricted Motif

	Amino Acid Position 1 2 3 4 5 6 7 8 9	Point Assignment
Dominant Binding Anchor Residue	L V	+3
Strong Binding Residue	M E V K K	+2
Weak Binding Residue	I A G I I A E L L Y P K L Y S F F D Y T H K P T N M M G Y S V H	+1

The derived peptide motif as currently defined is not particularly stringent. Some HLA-A2.1 binding peptides do not contain both dominant anchor residues and the amino acids flanking the dominant anchor residues play major roles in allowing or disallowing binding. Not every peptide with the current described binding motif will bind, and some peptides without the motif will bind. However, the current motif is valid enough to allow identification of some peptides capable of binding. Of note, the current HLA-A2.1 motif places 6 amino acids

09167516-100698

between the dominant anchor amino acids at residues 2 and 9.

Following identification of peptide motifs within a HER-2/neu polypeptide, amino acid substitutions may be made conservatively or non-conservatively. The latter type of substitutions are intended to produce an improved polypeptide that is more potent and/or more broadly cross-reactive (MHC polymorphism). An example of a more potent polypeptide is one that binds with higher affinity to the same MHC molecule as natural polypeptide, without affecting recognition by T cells specific for natural polypeptide. An example of a polypeptide with broader cross-reactivity is one that induces more broadly cross-reactive immune responses (i.e., binds to a greater range of MHC molecules) than natural polypeptide. Similarly, one or more amino acids residing between peptide motifs and having a spacer function (e.g., do not interact with a MHC molecule or T cell receptor) may be substituted conservatively or non-conservatively. It will be evident to those of ordinary skill in the art that polypeptides containing one or more amino acid substitutions may be tested for beneficial or adverse immunological interactions by a variety of assays, including those described herein for the ability to stimulate T cell recognition.

Variants within the scope of this invention may also, or alternatively, contain other modifications, including the deletion or addition of amino acids, that have minimal influence on the desired immunological properties of the polypeptide. It will be appreciated by those of ordinary skill in the art that truncated forms or non-native extended forms of a HER-2/neu polypeptide may be used, provided the desired immunological properties are at least roughly equivalent to that of full length, native

00167515-100698

HER-2/neu polypeptide. Cysteine residues may be deleted or replaced with other amino acids to prevent formation of incorrect intramolecular disulfide bridges upon renaturation. Other approaches to mutagenesis involve
5 modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present.

A HER-2/neu polypeptide may generally be obtained using a genomic or cDNA clone encoding the
10 protein. A genomic sequence that encodes full length HER-2/neu is shown in SEQ ID NO:1, and the deduced amino acid sequence is presented in SEQ ID NO:2. Such clones may be isolated by screening an appropriate expression library for clones that express HER-2/neu protein. The
15 library preparation and screen may generally be performed using methods known to those of ordinary skill in the art, such as methods described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, N.Y., 1989, which is
20 incorporated herein by reference. Briefly, a bacteriophage expression library may be plated and transferred to filters. The filters may then be incubated with a detection reagent. In the context of this invention, a "detection reagent" is any compound capable
25 of binding to HER-2/neu protein, which may then be detected by any of a variety of means known to those of ordinary skill in the art. Typical detection reagents contain a "binding agent," such as Protein A, Protein G, IgG or a lectin, coupled to a reporter group. Preferred
30 reporter groups include enzymes, substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. More preferably, the reporter group is horseradish peroxidase, which may be detected by incubation with a substrate such as

09167516.100698

09167516.100698

tetramethylbenzidine or 2,2'-azino-di-3-ethylbenz-thiazoline sulfonic acid. Plaques containing genomic or cDNA sequences that express HER-2/*neu* protein are isolated and purified by techniques known to those of ordinary skill in the art. Appropriate methods may be found, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, N.Y., 1989.

Variants of the polypeptide that retain the ability to stimulate an immune response may generally be identified by modifying the sequence in one or more of the aspects described above and assaying the resulting polypeptide for the ability to stimulate an immune response, e.g., a T cell response. For example, such assays may generally be performed by contacting T cells with the modified polypeptide and assaying the response. Naturally occurring variants of the polypeptide may also be isolated by, for example, screening an appropriate cDNA or genomic library with a DNA sequence encoding the polypeptide or a variant thereof.

The above-described sequence modifications may be introduced using standard recombinant techniques or by automated synthesis of the modified polypeptide. For example, mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analogue having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide a gene in which particular codons are altered according to the substitution, deletion, or insertion required.

Exemplary methods of making the alterations set forth above are disclosed by Walder et al., *Gene* 42:133, 1986; Bauer et al., *Gene* 37:73, 1985; Craik, *BioTechniques*, January 1985, 12-19; Smith et al., *Genetic Engineering: Principles and Methods*, Plenum Press, 1981; and U.S. Patent Nos. 4,518,584 and 4,737,462.

Mutations in nucleotide sequences constructed for expression of such HER-2/*neu* polypeptides must, of course, preserve the reading frame of the coding sequences and preferably will not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, which would adversely affect translation of the mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation *per se* be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the expressed HER-2/*neu* polypeptide mutants screened for the desired activity.

Not all mutations in a nucleotide sequence which encodes a HER-2/*neu* polypeptide will be expressed in the final product. For example, nucleotide substitutions may be made to enhance expression, primarily to avoid secondary structure loops in the transcribed mRNA (see, e.g., European Patent Application 75,444A), or to provide codons that are more readily translated by the selected host, such as the well-known *E. coli* preference codons for *E. coli* expression.

The polypeptides of the present invention, both naturally occurring and modified, are preferably produced by recombinant DNA methods. Such methods include inserting a DNA sequence encoding a HER-2/*neu* polypeptide into a recombinant expression vector and expressing the DNA sequence in a recombinant microbial, mammalian or

00167516-100690

insect cell expression system under conditions promoting expression. DNA sequences encoding the polypeptides provided by this invention can be assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being inserted in a recombinant expression vector and expressed in a recombinant transcriptional unit.

Recombinant expression vectors contain a DNA sequence encoding a HER-2/*neu* polypeptide operably linked to suitable transcriptional or translational regulatory elements derived from mammalian, microbial, viral or insect genes. Such regulatory elements include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and sequences which control the termination of transcription and translation. An origin of replication and a selectable marker to facilitate recognition of transformants may additionally be incorporated.

DNA regions are operably linked when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous and, in the case of secretory leaders, in reading frame. DNA sequences encoding HER-2/*neu* polypeptides which are to be expressed in a microorganism will preferably contain no introns that

00167516-100500

could prematurely terminate transcription of DNA into mRNA.

Expression vectors for bacterial use may comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. *E. coli* is typically transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar et al., *Gene* 2:95, 1977). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells.

Promoters commonly used in recombinant microbial expression vectors include the β -lactamase (penicillinase) and lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), the tryptophan (*trp*) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and European Patent Application 36,776) and the *tac* promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p.412, 1982). A particularly useful bacterial expression system employs the phage λ P_L promoter and *cI857ts* thermolabile repressor. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2, resident in *E. coli* strain JMB9 (ATCC 37092) and pPLc28, resident in *E. coli* RR1 (ATCC 53082).

Suitable promoter sequences in yeast vectors include the promoters for metallothionein, 3-

09157516.100608

phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-
5 3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Suitable vectors and promoters for use in
10 yeast expression are further described in R. Hitzeman et al., European Patent Application 73,657.

Preferred yeast vectors can be assembled using DNA sequences from pBR322 for selection and replication in *E. coli* (Amp^r gene and origin of replication) and yeast DNA
15 sequences including a glucose-repressible ADH2 promoter and α -factor secretion leader. The ADH2 promoter has been described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). The yeast α -factor leader, which directs secretion of heterologous
20 proteins, can be inserted between the promoter and the structural gene to be expressed (see, e.g., Kurjan et al., *Cell* 30:933, 1982; and Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984). The leader sequence may be modified to contain, near its 3' end, one or more useful
25 restriction sites to facilitate fusion of the leader sequence to foreign genes. The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters
30 and enhancers are derived from polyoma, adenovirus 2, simian virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other

genetic elements required for expression of a heterologous DNA sequence. The early and late promoters are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* II site located in the viral origin of replication is included. Further, viral genomic promoter, control and/or signal sequences may be utilized, provided such control sequences are compatible with the host cell chosen. Exemplary vectors can be constructed as disclosed by Okayama and Berg, *Mol. Cell. Biol.* 3:280, 1983.

15 A useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986). A preferred eukaryotic vector for expression of HER-2/neu polypeptide DNA is pDC406 (McMahan et al., *EMBO J.* 10:2821, 1991), and includes regulatory sequences derived from SV40, human immunodeficiency virus (HIV), and Epstein-Barr virus (EBV). Other preferred vectors include pDC409 and pDC410, which are derived from pDC406. pDC410 was derived from pDC406 by substituting the EBV origin of replication with sequences encoding the SV40 large T antigen. pDC409 differs from pDC406 in that a *Bgl* II restriction site outside of the multiple cloning site has been deleted, making the *Bgl* II site within the multiple cloning site unique.

30 A useful cell line that allows for episomal replication of expression vectors, such as pDC406 and pDC409, which contain the EBV origin of replication, is CV-1/EBNA (ATCC CRL 10478). The CV-L/EBNA cell line was

derived by transfection of the CV-1 cell line with a gene encoding Epstein-Barr virus nuclear antigen-I (EBNA-1) and constitutively express EBNA-1 driven from human CMV immediate-early enhancer/promoter.

5 Transformed host cells are cells which have been transformed or transfected with expression vectors constructed using recombinant DNA techniques and which contain sequences encoding a HER-2/*neu* polypeptide of the present invention. Transformed host cells may express the
10 desired HER-2/*neu* polypeptide, but host cells transformed for purposes of cloning or amplifying HER-2/*neu* DNA do not need to express the HER-2/*neu* polypeptide. Expressed polypeptides will preferably be secreted into the culture supernatant, depending on the DNA selected, but may also
15 be deposited in the cell membrane.

Suitable host cells for expression of recombinant proteins include prokaryotes, yeast or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram
20 positive organisms, for example *E. coli* or *Bacilli*. Higher eukaryotic cells include established cell lines of insect or mammalian origin as described below. Cell-free translation systems could also be employed to produce HER-2/*neu* polypeptides using RNAs derived from DNA
25 constructs. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, by Pouwels et al., *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985.

30 Prokaryotic expression hosts may be used for expression of HER-2/*neu* polypeptides that do not require extensive proteolytic and disulfide processing. Prokaryotic expression vectors generally comprise one or more phenotypic selectable markers, for example a gene

09167516.106360

encoding proteins conferring antibiotic resistance or supplying an autotrophic requirement, and an origin of replication recognized by the host to ensure amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although other hosts may also be employed.

Recombinant HER-2/neu polypeptides may also be expressed in yeast hosts, preferably from the *Saccharomyces* species, such as *S. cerevisiae*. Yeast of other genera, such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will generally contain an origin of replication from the 2 μ yeast plasmid or an autonomously replicating sequence (ARS), a promoter, DNA encoding the HER-2/neu polypeptide, sequences for polyadenylation and transcription termination and a selection gene. Preferably, yeast vectors will include an origin of replication and selectable marker permitting transformation of both yeast and *E. coli*, e.g., the ampicillin resistance gene of *E. coli* and the *S. cerevisiae* *trp1* gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, and a promoter derived from a highly expressed yeast gene to induce transcription of a structural sequence downstream. The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

Suitable yeast transformation protocols are known to those of skill in the art. An exemplary technique described by Hind et al. (*Proc. Natl. Acad. Sci. USA* 75:1929, 1978), involves selecting for Trp⁺ transformants in a selective medium consisting of 0.67%

yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 mg/ml adenine and 20 mg/ml uracil. Host strains transformed by vectors comprising the ADH2 promoter may be grown for expression in a rich medium consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80 mg/ml adenine and 80 mg/ml uracil. Derepression of the ADH2 promoter occurs upon exhaustion of medium glucose. Crude yeast supernatants are harvested by filtration and held at 4°C prior to further purification.

10 Various mammalian or insect (e.g., *Spodoptera* or *Trichoplusia*) cell culture systems can also be employed to express recombinant polypeptide. Baculovirus systems for production of heterologous polypeptides in insect cells are reviewed, for example, by Luckow and Summers, 15 *Bio/Technology* 6:47, 1988. Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (*Cell* 23:175, 1981), and other cell lines capable of expressing an appropriate vector including, for example, CV-1/EBNA (ATCC CRL 10478), L 20 cells, C127, 3T3, Chinese hamster ovary (CHO), COS, NS-1, HeLa and BHK cell lines. Mammalian expression vectors may comprise nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking 25 nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

30 Purified HER-2/*neu* polypeptides may be prepared by culturing suitable host/vector systems to express the recombinant translation products of the DNAs of the present invention, which are then purified from culture media or cell extracts. For example, supernatants from systems which secrete recombinant polypeptide into culture

00167515-100600

media may be first concentrated using a commercially available protein concentration filter, such as an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate may be applied to a suitable purification matrix. For example, a suitable affinity matrix may comprise a counter structure protein (i.e., a protein to which a HER-2/neu polypeptide binds in a specific interaction based on structure) or lectin or antibody molecule bound to a suitable support.

- 10 Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, 15 a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred. Gel filtration chromatography also provides a means of purifying a HER-2/neu.

- 20 Affinity chromatography is a preferred method of purifying HER-2/neu polypeptides. For example, monoclonal antibodies against the HER-2/neu polypeptide may also be useful in affinity chromatography purification, by utilizing methods that are well-known in the art.

- 25 Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media (e.g., silica gel having pendant methyl or other aliphatic groups) may be employed to further purify a HER-2/neu polypeptide 30 composition. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a homogeneous recombinant polypeptide.

Recombinant HER-2/neu polypeptide produced in bacterial culture is preferably isolated by initial

09167516-100698

extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange or size exclusion chromatography steps. High performance liquid chromatography (HPLC) may be employed for final
5 purification steps. Microbial cells employed in expression of recombinant HER-2/neu polypeptide can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

10 Fermentation of yeast which express HER-2/neu polypeptide as a secreted protein greatly simplifies purification. Secreted recombinant protein resulting from a large-scale fermentation can be purified by methods analogous to those disclosed by Urdal et al. (J.
15 Chromatog. 296:171, 1984). This reference describes two sequential, reverse-phase HPLC steps for purification of recombinant human GM-CSF on a preparative HPLC column.

Preparations of HER-2/neu polypeptides synthesized in recombinant culture may contain non-
20 HER-2/neu cell components, including proteins, in amounts and of a character which depend upon the purification steps taken to recover the HER-2/neu polypeptide from the culture. These components ordinarily will be of yeast, prokaryotic or non-human eukaryotic origin. Such
25 preparations are typically free of other proteins which may be normally associated with the HER-2/neu protein as it is found in nature in its species of origin.

Automated synthesis provides an alternate method for preparing polypeptides of this invention. For
30 example, any of the commercially available solid-phase techniques may be employed, such as the Merrifield solid phase synthesis method, in which amino acids are sequentially added to a growing amino acid chain. (See Merrifield, J. Am. Chem. Soc. 85:2149-2146, 1963.)

09167516.100698

Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied Biosystems, Inc. of Foster City, CA, and may generally be operated according to the manufacturer's instructions.

- 5 Within one aspect of the present invention, use of a HER-2/neu polypeptide (or a DNA molecule that directs the expression of such a peptide) to generate an immune response to the HER-2/neu protein (including that expressed on a malignancy in which a HER-2/neu oncogene is associated) may be detected. Representative examples of such malignancies include breast, ovarian, colon, lung and prostate cancers. An immune response to the HER-2/neu protein, once generated by a HER-2/neu polypeptide, can be long-lived and can be detected long after immunization, 10 regardless of whether the protein is present or absent in the body at the time of testing. An immune response to the HER-2/neu protein generated by reaction to a HER-2/neu polypeptide can be detected by examining for the presence or absence, or enhancement, of specific activation of CD4⁺ 15 or CD8⁺ T cells. More specifically, T cells isolated from an immunized individual by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes) are incubated with HER-2/neu protein. For example, T cells may be incubated *in vitro* 20 for 2-9 days (typically 4 days) at 37°C with HER-2/neu protein (typically, 5 µg/ml of whole protein or graded numbers of cells synthesizing HER-2/neu protein). It may be desirable to incubate another aliquot of a T cell sample in the absence of HER-2/neu protein to serve as a 25 control.

Specific activation of CD4⁺ or CD8⁺ T cells may be detected in a variety of ways. Methods for detecting specific T cell activation include detecting the proliferation of T cells, the production of cytokines

00167516-100699

(e.g., lymphokines), or the generation of cytolytic activity (i.e., generation of cytotoxic T cells specific for HER-2/neu protein). For CD4⁺ T cells, a preferred method for detecting specific T cell activation is the
5 detection of the proliferation of T cells. For CD8⁺ T cells, a preferred method for detecting specific T cell activation is the detection of the generation of cytolytic activity.

Detection of the proliferation of T cells may be
10 accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring the rate of DNA synthesis. T cells which have been stimulated to proliferate exhibit an increased rate of DNA synthesis. A typical way to measure the rate of DNA
15 synthesis is, for example, by pulse-labeling cultures of T cells with tritiated thymidine, a nucleoside precursor which is incorporated into newly synthesized DNA. The amount of tritiated thymidine incorporated can be determined using a liquid scintillation spectrophotometer.
20 Other ways to detect T cell proliferation include measuring increases in interleukin-2 (IL-2) production, Ca²⁺ flux, or dye uptake, such as 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium. Alternatively, synthesis of lymphokines (such as interferon-gamma) can be measured or
25 the relative number of T cells that can respond to intact p185^{HER-2/neu} protein may be quantified.

By use or expression of a HER-2/neu polypeptide, T cells which recognize the HER-2/neu protein can be proliferated *in vivo*. For example, immunization with a
30 HER-2/neu peptide (i.e., as a vaccine) can induce continued expansion in the number of T cells necessary for therapeutic attack against a tumor in which the HER-2/neu oncogene is associated. Typically, about 0.01 µg/kg to about 100 mg/kg body weight will be administered by the

09167516-100698

intradermal, subcutaneous or intravenous route. A preferred dosage is about 1 µg/kg to about 1 mg/kg, with about 5 µg/kg to about 200 µg/kg particularly preferred. It will be evident to those skilled in the art that the number and frequency of administration will be dependent upon the response of the patient. It may be desirable to administer the HER-2/neu polypeptide repetitively. It will be evident to those skilled in this art that more than one HER-2/neu polypeptide may be administered, either simultaneously or sequentially. Preferred peptides for immunization are those that include the amino acid sequence of SEQ ID NO:2 beginning at about the lysine residue at amino acid position 676 and extending to about the valine residue at amino acid position 1255. It will be appreciated by those in the art that the present invention contemplates the use of an intact HER-2/neu polypeptide as well as division of such a polypeptide into a plurality of peptides. Neither intact p185^{HER-2/neu} protein nor a peptide having the amino acid sequence of its entire extracellular domain (i.e., a peptide having an amino acid sequence of SEQ ID NO:2 from amino acid position 1 up to amino acid position 650, plus or minus about one to five positions, and with or without the first 21 amino acid positions) are used alone for immunization.

A HER-2/neu polypeptide (or nucleic acid) is preferably formulated for use in the above methods as a pharmaceutical composition (e.g., vaccine). Pharmaceutical compositions generally comprise one or more polypeptides in combination with a pharmaceutically acceptable carrier, excipient or diluent. Such carriers will be nontoxic to recipients at the dosages and concentrations employed. The use of a HER-2/neu polypeptide in conjunction with chemotherapeutic agents is also contemplated.

09167515.100698

09157516-100608

In addition to the HER-2/*neu* polypeptide (which functions as an antigen), it may be desirable to include other components in the vaccine, such as a vehicle for antigen delivery and immunostimulatory substances designed to enhance the protein's immunogenicity. Examples of vehicles for antigen delivery include aluminum salts, water-in-oil emulsions, biodegradable oil vehicles, oil-in-water emulsions, biodegradable microcapsules, and liposomes. Examples of immunostimulatory substances (adjuvants) include N-acetylmuramyl-L-alanine-D-isoglutamine (MDP), lipopoly-saccharides (LPS), glucan, IL-12, GM-CSF, gamma interferon and IL-15. It will be evident to those of ordinary skill in this art that a HER-2/*neu* polypeptide for a vaccine may be prepared synthetically or be naturally derived.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration and whether a sustained release is desired. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (e.g., polylactic galactide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109. A HER-2/*neu* polypeptide may be encapsulated within the biodegradable microsphere or associated with the surface of the microsphere. For example, in a

preferred embodiment, a polypeptide having the amino acid sequence of SEQ ID NO:2 from amino acid 676 through amino acid 1255 is encapsulated within a biodegradable microsphere. In this regard, it is preferable that the microsphere be larger than approximately 25 microns.

Pharmaceutical compositions (including vaccines) may also contain diluents such as buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with nonspecific serum albumin are exemplary appropriate diluents. Preferably, product is formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents.

As an alternative to the presentation of HER-2/neu polypeptides, the subject invention includes compositions capable of delivering nucleic acid molecules encoding a HER-2/neu polypeptide. Such compositions include recombinant viral vectors (e.g., retroviruses (see WO 90/07936, WO 91/02805, WO 93/25234, WO 93/25698, and WO 94/03622), adenovirus (see Berkner, *Biotechniques* 6:616-627, 1988; Li et al., *Hum. Gene Ther.* 4:403-409, 1993; Vincent et al., *Nat. Genet.* 5:130-134, 1993; and Kolls et al., *Proc. Natl. Acad. Sci. USA* 91:215-219, 1994), pox virus (see U.S. Patent No. 4,769,330; U.S. Patent No. 5,017,487; and WO 89/01973)), naked DNA (see WO 90/11092), nucleic acid molecule complexed to a polycationic molecule (see WO 93/03709), and nucleic acid associated with liposomes (see Wang et al., *Proc. Natl. Acad. Sci. USA* 84:7851, 1987). In certain embodiments, the DNA may be linked to killed or inactivated adenovirus (see Curiel et al., *Hum. Gene Ther.* 3:147-154, 1992; Cotton et al., *Proc.*

09167516.100698

Natl. Acad. Sci. USA 89:6094, 1992). Other suitable compositions include DNA-ligand (see Wu et al., *J. Biol. Chem.* 264:16985-16987, 1989) and lipid-DNA combinations (see Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1989). In addition, the efficiency of naked DNA uptake into cells may be increased by coating the DNA onto biodegradable beads.

In addition to direct *in vivo* procedures, *ex vivo* procedures may be used in which cells are removed from an animal, modified, and placed into the same or another animal. It will be evident that one can utilize any of the compositions noted above for introduction of HER-2/*neu* nucleic acid molecules into tissue cells in an *ex vivo* context. Protocols for viral, physical and chemical methods of uptake are well known in the art.

Accordingly, the present invention is useful for enhancing or eliciting, in a patient or cell culture, a cellular immune response (e.g., the generation of antigen-specific cytolytic T cells). As used herein, the term "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with cancer, such as breast cancer, or may be normal (i.e., free of detectable disease and infection). A "cell culture" is any preparation of T cells or isolated component cells (including, but not limited to, macrophages, monocytes, B cells and dendritic cells). Such cells may be isolated by any of a variety of techniques well known to those of ordinary skill in the art (such as Ficoll-hypaque density centrifugation). The cells may (but need not) have been isolated from a patient afflicted with a HER-2/*neu* associated malignancy, and may be reintroduced into a patient after treatment.

The present invention also discloses that HER-2/*neu* polypeptide, in addition to being immunogenic to

09167516.100698

09167316-100608

T cells, appears to stimulate B-cells to produce antibodies capable of recognizing HER-2/*neu* polypeptide. Antibodies specific (i.e., which exhibit a binding affinity of about 10^7 liters/mole or better) for HER-2/*neu* protein may be found in a variety of body fluids including sera and ascites. Briefly, a body fluid sample is isolated from a warm-blooded animal, such as a human, for whom it is desired to determine whether antibodies specific for HER-2/*neu* polypeptide are present. The body fluid is incubated with HER-2/*neu* polypeptide under conditions and for a time sufficient to permit immunocomplexes to form between the polypeptide and antibodies specific for the protein. For example, a body fluid and HER-2/*neu* polypeptide may be incubated at 4°C for 24-48 hours. Following the incubation, the reaction mixture is tested for the presence of immunocomplexes. Detection of one or more immunocomplexes formed between HER-2/*neu* polypeptide and antibodies specific for HER-2/*neu* polypeptide may be accomplished by a variety of known techniques, such as radioimmunoassays (RIA) and enzyme linked immunosorbent assays (ELISA).

Suitable immunoassays include the double monoclonal antibody sandwich immunoassay technique of David et al. (U.S. Patent 4,376,110); monoclonal polyclonal antibody sandwich assays (Wide et al., in Kirkham and Hunter, eds., *Radioimmunoassay Methods*, E. and S. Livingstone, Edinburgh, 1970); the "western blot" method of Gordon et al. (U.S. Patent 4,452,901); immunoprecipitation of labeled ligand (Brown et al., *J. Biol. Chem.* 255:4980-4983, 1980); enzyme-linked immunosorbent assays as described by, for example, Raines and Ross (*J. Biol. Chem.* 257:5154-5160, 1982); immunocytochemical techniques, including the use of fluorochromes (Brooks et al., *Clin. Exp. Immunol.* 39: 477,

1980); and neutralization of activity [Bowen-Pope et al., *Proc. Natl. Acad. Sci. USA* 81:2396-2400 (1984)], all of which are hereby incorporated by reference. In addition to the immunoassays described above, a number of other
5 immunoassays are available, including those described in U.S. Patent Nos.: 3,817,827; 3,850,752; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; and 4,098,876, all of which are herein incorporated by reference.

For detection purposes, HER-2/*neu* polypeptide
10 ("antigen") may either be labeled or unlabeled. When unlabeled, the antigen finds use in agglutination assays. In addition, unlabeled antigen can be used in combination with labeled molecules that are reactive with immunocomplexes, or in combination with labeled antibodies
15 (second antibodies) that are reactive with the antibody directed against HER-2/*neu* polypeptide, such as antibodies specific for immunoglobulin. Alternatively, the antigen can be directly labeled. Where it is labeled, the reporter group can include radioisotopes, fluorophores,
20 enzymes, luminescers, or dye particles. These and other labels are well known in the art and are described, for example, in the following U.S. patents: 3,766,162; 3,791,932; 3,817,837; 3,996,345; and 4,233,402.

Typically in an ELISA assay, antigen is adsorbed
25 to the surface of a microtiter well. Residual protein-binding sites on the surface are then blocked with an appropriate agent, such as bovine serum albumin (BSA), heat-inactivated normal goat serum (NGS), or BLOTTO (buffered solution of nonfat dry milk which also contains
30 a preservative, salts, and an antifoaming agent). The well is then incubated with a sample suspected of containing specific antibody. The sample can be applied neat, or, more often, it can be diluted, usually in a buffered solution which contains a small amount (0.1%-5.0%

09167516-100698

by weight) of protein, such as BSA, NGS, or BLOTTO. After incubating for a sufficient length of time to allow specific binding to occur, the well is washed to remove unbound protein and then incubated with an anti-species
5 specific immunoglobulin antibody labeled with a reporter group. The reporter group can be chosen from a variety of enzymes, including horseradish peroxidase, beta-galactosidase, alkaline phosphatase, and glucose oxidase. Sufficient time is allowed for specific binding to occur,
10 then the well is again washed to remove unbound conjugate, and the substrate for the enzyme is added. Color is allowed to develop and the optical density of the contents of the well is determined visually or instrumentally.

In one preferred embodiment of this aspect of
15 the present invention, a reporter group is bound to HER-2/neu protein. The step of detecting immunocomplexes involves removing substantially any unbound HER-2/neu protein and then detecting the presence or absence of the reporter group.

20 In another preferred embodiment, a reporter group is bound to a second antibody capable of binding to the antibodies specific for HER-2/neu protein. The step of detecting immunocomplexes involves (a) removing substantially any unbound antibody, (b) adding the second
25 antibody, (c) removing substantially any unbound second antibody and then (d) detecting the presence or absence of the reporter group. Where the antibody specific for HER-2/neu protein is derived from a human, the second antibody is an anti-human antibody.

30 In a third preferred embodiment for detecting immunocomplexes, a reporter group is bound to a molecule capable of binding to the immunocomplexes. The step of detecting involves (a) adding the molecule, (b) removing substantially any unbound molecule, and then (c) detecting

09167516.100698

the presence or absence of the reporter group. An example of a molecule capable of binding to the immunocomplexes is protein A.

It will be evident to one skilled in the art that a variety of methods for detecting the immunocomplexes may be employed within the present invention. Reporter groups suitable for use in any of the methods include radioisotopes, fluorophores, enzymes, luminescers, and dye particles.

In a related aspect of the present invention, detection of immunocomplexes formed between HER-2/*neu* polypeptide and antibodies in body fluid which are specific for HER-2/*neu* polypeptide may be used to monitor the effectiveness of cancer therapy, which involves a HER-2/*neu* polypeptide, for a malignancy in which the HER-2/*neu* oncogene is associated. Samples of body fluid taken from an individual prior to and subsequent to initiation of therapy may be analyzed for the immunocomplexes by the methodologies described above. Briefly, the number of immunocomplexes detected in both samples are compared. A substantial change in the number of immunocomplexes in the second sample (post-therapy initiation) relative to the first sample (pre-therapy) reflects successful therapy.

The following examples are offered by way of illustration and not by way of limitation.

00167515-100559

EXAMPLES

EXAMPLE 1

5 EXPRESSION AND PURIFICATION OF RECOMBINANT HUMAN HER-2/*neu* POLYPEPTIDE

09167516.100698

The human HER-2/*neu* polypeptide was recovered by the PCR method (e.g., U.S. Patent Nos. 4,683,195; 4,683,202; 4,800,159) from a plasmid prepared according to Di Fiore et al. (King et al., *Science* 229:974-976, 1985; Di Fiore et al., *Science* 237:178-182, 1987) using oligonucleotide primers that additionally introduced a BssHII restriction site and an enterokinase protease site on the 5' end and an EcoRI site on the 3' end. The primer 15 for the 5'-end was 5'-TCTGGCGCGCTGGATGACGATGACAAGAAACGACGGCAGCAGAAGATC-3' (SEQ ID NO:3) while the primer for the 3'-end was 5'-TGAATTCTCGAGTCATTACACTGGCAGCTCCAGACCCAG-3' (SEQ ID NO:4). The resulting 1.8 kb PCR fragment was subcloned 20 into the T-vector from Novagen (Madison, WI, USA) and the sequence of selected clones was determined on the ABI 373 automated DNA sequencer (Applied Biosystems Inc., Foster City, CA, USA) using overlapping sequencing primers. PCR fragments with sequence that corresponded to the published 25 DNA sequence for the human HER-2/*neu* cDNA (SEQ ID NO:1; Coussens et al., *Science* 230:1132, 1985; Yamamoto et al., *Nature* 319:230, 1986) were then connected in the correct reading frame via the BssHII site to a modified *E. coli* thioredoxin reductase. A 6Xhistidine affinity tag 30 employed in Ni-NTA affinity purification of the expressed fusion protein was incorporated into the thioredoxin reductase fusion partner. This cDNA for the *trx*A-human HER-2/*neu* polypeptide fusion protein was subcloned into a modified pET expression vector for expression in *E. coli*.

While thioredoxin reductase has been reported to stabilize and solubilize other heterologous proteins expressed in *E. coli*, it did not appear to offer any significant advantage for human HER-2/*neu* polypeptide expression in *E. coli*. While a significant proportion of the trxA-HER-2/*neu* polypeptide fusion protein was soluble, a majority was expressed in inclusion bodies. The fusion protein was also subjected to degradation during expression in *E. coli*. The presence of the thioredoxin reductase fusion partner may, however, stabilize the protein during purification. The availability of monoclonal antibodies to thioredoxin reductase provides a convenient marker to follow during purification.

For purification of the human HER-2/*neu* polypeptide with the thioredoxin reductase fusion partner containing the 6XHis affinity tag, the *E. coli* pellet was resuspended with protease inhibitors and lysozyme and sonicated. The inclusion bodies were isolated by centrifugation, and are washed 3X with deoxycholate, the last wash being overnight to remove LPS. The washed inclusion bodies are solubilized in GuHCl for Ni purification. The Ni column was eluted with Imidazole in urea and dialyzed against 10 mM Tris pH8. The recovery of HER-2/*neu* polypeptide using this protocol was from 80%-95% pure full length protein with the main contaminant being degraded protein. From 500 ml of fermentation, 20 mg were recovered. It was >98% HER-2/*neu* polypeptide. The techniques used herein are well known to those in the art and have been described, for example, in J. Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, 1989, Cold Spring Harbor, New York, USA.

EXAMPLE 2

DENDRITIC CELLS CAN PRIME HUMAN HER-2/NEU POLYPEPTIDE

A. Generation of DC Cultures From Bone Marrow

- 5 DC cultures were generated from CD34+ hematopoietic progenitor cells (HPC). CD34+ cells were purified from bone marrow of normal donors using the cell separation system Ceparate LC Kit (CellPro, Bothell, WA, USA). Purity of recovered CD34+ cells was determined by
- 10 flow cytometric analysis to be 80% to 95%. CD34+ cells were cultured in serum-free medium (X-VIVO 10, Biowhittaker, Inc., Walkersville, MD, USA) supplemented with L-glutamine (584 µg/l), penicillin (10 IU/ml), streptomycin (100 µg/ml), 100 ng/ml human rGM-CSF and 50
- 15 ng/ml human rIL-6 (Immunex, Seattle, WA, USA). After 0 to 17 days of culture time, cells were harvested and used for phenotyping and T cell stimulation assays. GM-CSF alone and in combination with IL-4 or TNFα have been described to induce the *in vitro* growth of DC. In experiments using
- 20 KLH and OVA as antigens to prime naive T cells, GM-CSF plus IL-6 consistently gave a comparable total stimulation, but with a lower background and thus a higher stimulation index as compared to GM-CSF plus IL-4 or TNFα.

25 B. T Cell Priming Assay

- Bone marrow derived CD34+ HPC cultured in serum-free medium containing GM-CSF and IL-6 were used as APC after a culture period of 0-17 days. Priming ability of DC was determined by culturing them with autologous, naive
- 30 T lymphocytes in the presence or absence of the protein antigen recombinant human HER-2/neu polypeptide (hHNP) (10 µg/ml). CD4+ T lymphocytes were isolated from peripheral blood mononuclear cells by positive selection using

09167516-100698

immunoaffinity columns (CellPro, Inc., Bothell, WA, USA). CD4+ CD45RA+ (naive) T lymphocytes were selected from CD4+ T lymphocytes using an anti-CD45RA mAb directly conjugated to FITC (Immunotech, Westbrook, ME, USA) by flow cytometric sorting. The CD4+ CD45RA+ T cells obtained were 99% pure. DC cultures were plated into 96-well round-bottomed plates (Corning, Corning, NY, USA) at various concentrations and incubated for 16-18 hours with hHNP 10 $\mu\text{g/ml}$ final concentration. Antigen-pulsed DC were irradiated (10 Gy), and autologous CD4+ CD45RA+ T lymphocytes were added ($5 \times 10^4/\text{well}$). Proliferative response of T cells was measured by the uptake of (^3H)thymidine (1 $\mu\text{Ci/well}$) added on day 6 for 16-18 hours. Proliferation assays were performed in serum-free and cytokine-free medium. The results are shown in Figure 1. Figure 2 shows the results of testing CD4+ T cells, from a normal donor, for responses to hHNP. Similar data was obtained with T cells from nine out of ten normal individuals.

20

EXAMPLE 3

ASSAY FOR DETECTING LOW FREQUENCY LYMPHOCYTE PRECURSORS

Three assays can be used for the detection of CD4+ responses: a standard proliferation assay, a screening method for low frequency events, and a limiting dilution assay (LDA). Conventional proliferative assays are capable of readily detecting primed responses. The proliferative response stimulation index provides a rough correlation with precursor frequency of antigen-reactive T cells. Any specific proliferative response detected from PBL is considered to be a primed response.

30

09167516-100698

To provide a more quantitative interpretation of CD4⁺ T cell responses, the assay system developed for detecting low lymphocyte precursor frequency responses (described below) is used. This assay is simple and cost-effective. In circumstances in which more precision is needed, the precursor frequency is validated by limiting dilution assays (Bishop and Orosz, *Transplantation* 47:671-677, 1989).

Responses greater than detected in normal individuals are defined as a primed response and imply existent immunity. Low responses, detectable only by LDA conditions are considered to be unprimed responses. An absent response by LDA or a response lower than that defined by the normal population analysis is considered to be tolerance/anergy.

In general, primed CD4⁺ T cell responses can be detected in conventional proliferative assays, whereas unprimed responses are not detectable in the same assays. Detection of small numbers of unprimed T cells is limited by confounding background thymidine uptake including the autologous mixed lymphocyte response (AMLR) to self MHC antigen plus responses to processed self serum proteins and exogenously added serum proteins.

To elicit and detect unprimed T cells, an assay system for low frequency responses based on Poisson sampling statistics was used (In: *Pinnacles*, Chiron Corporation, 1:1-2, 1991). This type of analysis applies specifically to low frequency events in that, if the precursor frequency is less than the number of cells in one replicate culture, many replicates are required to detect a statistically significant number of positives. Theoretically, the analysis will correct for autologous responses by setting up a known positive control (such as PHA or tetanus toxoid) and known negative control (no

antigen) and evaluating all data points from lowest to highest irrespective of the experimental group to which they belong. A cutoff value is calculated based on the equation $\text{cutoff} = M + (F + SD)$, where M = arithmetic mean, $F = 3.29$, a factor from tables of standardized normal distribution chosen so not more than 0.1% of the "true negatives" of a normally distributed background will be above the cutoff, and SD = standard deviation. In this screening assay, wells above the cutoff are considered true positives that potentially contain a lymphocyte that is specifically proliferating to the antigen of interest. Although estimations of lymphocyte precursor frequency is possible using this method, precise determination requires formal LDA analysis.

EXAMPLE 4

HER-2/NEU POLYPEPTIDE BASED VACCINE ELICITS IMMUNITY TO HER-2/NEU PROTEIN

A. Animals

Rats used in this study were Fischer strain 344 (CDF (F-344)/Cr1BR) (Charles River Laboratories, Portage MI). Animals were maintained at the University of Washington Animal facilities under specific pathogen free conditions and routinely used for experimental studies between 3 and 4 months of age.

B. Immunization

Fischer rats were immunized with recombinant rat HER-2/neu polypeptide (rHNP) in a variety of adjuvants (MPL, Vaxcel; Rib, Bozeman, MT, USA). Animals received 50 μg of rHNP mixed with adjuvant subcutaneously. Twenty days later the animals were boosted with a second

immunization of 50 μ g of rHNP administered in the same fashion. Twenty days after the booster immunization animals were tested for the presence of antibodies directed against rat HER-2/*neu* protein (*neu*).

5

C. Cell Lines

Two cell lines were used as a source of *neu* proteins. SKBR3, a human breast cancer cell line that is a marked overexpressor of HER-2/*neu* (American Type Culture Collection, Rockville, MD), was maintained in culture in 10% fetal bovine serum (FBS) (Gemini Bioproducts, Inc., Calabasas, CA) and RPMI. DHFR-G8, an NIH/3T3 cell line cotransfected with *cneu-p* and pSV2-DHFR (American Type Culture Collection, Rockville, MD), was used as a source of non-transforming rat *neu* protein (Bernards et al., *Proc. Natl. Acad. Sci. USA* 84:6854-6858, 1987). This cell line was maintained in 10% FBS and Dulbecco's modified Eagle's medium with 4.5g/L glucose. DHFR-G8 cells were passaged through the same medium supplemented with 0.3 μ M methotrexate at every third passage to maintain the *neu* transfectant.

D. Preparation of Cell Lysates

Lysates of both SKBR3 and DHFR-G8 were prepared and used as a source of *neu* protein. Briefly, a lysis buffer consisting of tris base, sodium chloride and Triton-X (1%) pH 7.5 was prepared. Protease inhibitors were added; aprotinin (1 μ g/ml), benzamidine (1mM) and PMSF (1mM). 1 ml of the lysis buffer was used to suspend 10⁷ cells. The cells were vortexed for 15 seconds every 10 minutes for an hour until disrupted. All procedures were performed on ice in a 4°C cold room. After disruption the cells were microfuged at 4°C for 20 minutes. Supernatant was removed from cell debris and stored in small aliquots

09167516-100698

at -70°C until used. Presence of human and rat *neu* in the lysates was documented by Western blot analysis.

E. ELISA for Rat *neu* Antibody Responses

5 96 well Immulon 4 plates (Baxter SP, Redmond, WA: Dynatech Laboratories) were incubated overnight at 4°C with a rat *neu* specific monoclonal antibody (Oncogene Science), 7.16.4, at a concentration of 10 µg/ml diluted in carbonate buffer (equimolar concentrations of Na₂CO₃ and
10 NaHCO₃ pH 9.6). After incubation, all wells were blocked with PBS-1% BSA (Sigma Chemical, St. Louis, MO, USA), 100 µl/well for 3 hours at room temperature. The plate was washed with PBS-0.5% Tween and lysates of DHFRG8, a murine cell line transfected with rat *neu* DNA (American Type
15 Culture Collection, Rockville, MD, USA); a source of rat *neu* protein, were added to alternating rows. The plate was incubated overnight at 4°C. The plate was then washed with PBS-0.5% Tween and experimental sera was added at the following dilutions: 1:25 to 1:200. The sera was diluted
20 in PBS-1% BSA-1% FBS-25 µg/ml mouse IgG-0.01% NaN₃ and then serially into PBS-1% BSA. 50 µl of diluted sera was added/well and incubated 1 hour at room temperature. Each experimental sera was added to a well with rat *neu* and a well without rat *neu*. Sheep anti-rat Ig F(ab')₂
25 horseradish peroxidase (HRP) was added to the wells at a 1:5000 dilution in PBS-1% BSA and incubated for 45 minutes at room temperature (Amersham Co., Arlington Heights, IL, USA). Following the final wash, TMB (Kirkegaard and Perry Laboratories, Gaithersburg, MD) developing reagent was
30 added. Color reaction was read at an optical density of 450 nm. The OD of each serum dilution was calculated as the OD of the rat *neu* coated wells minus the OD of the PBS-1% BSA coated wells. Sera from animals immunized with

09167516-100698

the adjuvants alone and an animal immunized with hHNP (foreign protein) were also evaluated in a similar manner. The results are shown in Figure 3.

5 F. T Cell Proliferation Assays

For analysis of HER-2/*neu* polypeptide specific responses: Fresh spleen or lymph node cells are harvested by mechanical disruption and passage through wire mesh and washed. 2 x 10⁵ spleen cells/well and 1 x 10⁵ lymph node
10 cells/well are plated into 96-well round bottom microtiter plates (Corning, Corning, NY) with 6 replicates per experimental group. The media consists of EHAA 120 (Biofluids) with L-glutamine, penicillin/streptomycin, 2-mercaptoethanol, and 5% FBS. Cells are incubated with
15 polypeptides. After 4 days, wells are pulsed with 1 μ Ci of [³H]thymidine for 6-8 hours and counted. Data is expressed as a stimulation index (SI) which is defined as the mean of the experimental wells divided by the mean of the control wells (no antigen). For analysis of HER-2/*neu*
20 protein specific responses: Spleen or lymph node cells are cultured for 3 in vitro stimulations. At the time of analysis 1 x 10⁵ cultured spleen or lymph node T cells are plated into 96 well microtiter plates as described above. Cells are incubated with 1 μ g/ml immunoaffinity column
25 purified rat *neu* (from DHFR-G8 cells as the source of rat *neu*). After 4 days, wells were pulsed with 1 μ Ci of [³H]thymidine for 6-8 hours and counted. Data is expressed as a stimulation index which is defined as the mean of the experimental wells divided by the mean of the control
30 wells (no antigen).

09167516-100698

EXAMPLE 5

PRIMED RESPONSES TO HUMAN HER-2/*neu* POLYPEPTIDE
CAN BE DETECTED IN PATIENTS WITH BREAST CANCER

5 Heparinized blood was obtained from a patient
with stage II HER-2/*neu* overexpressing breast cancer.
Peripheral blood mononuclear cells (PBMC) were separated
by Ficoll Hypaque density centrifugation. PBMC were
10 plated at a concentration of 2×10^5 /well into 96-well
round-bottomed plates (Corning, Corning, NY, USA). 24
well replicates were performed for each experimental
group. Antigens consisting of HER-2/*neu* derived peptides
(15-20 amino acids in length with number of first amino
acid in sequence listed) 25 $\mu\text{g/ml}$, human HER-2/*neu*
15 polypeptide (hHNP) 1 $\mu\text{g/ml}$, tetanus toxoid 1 $\mu\text{g/ml}$, and
p30 a peptide derived from tetanus 25 $\mu\text{g/ml}$ were added to
each 24 well replicate. The assay was performed in media
containing 10% human sera. Proliferative response of T
cells was measured by the uptake of (^3H)thymidine
20 (1 $\mu\text{Ci/well}$) added on day 4 for 10 hours. Positive wells,
antigen reactive wells, were scored as positive if the cpm
was greater than the mean and 3 standard deviations of the
no antigen wells. The results are shown in Figure 4.
This stage II breast cancer patient has a significant
25 response to recombinant hHNP.

All publications and patent applications
mentioned in this specification are herein incorporated by
30 reference to the same extent as if each individual
publication or patent application was specifically and
individually incorporated by reference.

From the foregoing, it will be evident that,
although specific embodiments of the invention have been

00167516-100698

described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

09167516-100639
869001-91529160

Sequence Listing

(1) GENERAL INFORMATION:

(i) APPLICANT: University of Washington

(ii) TITLE OF INVENTION: COMPOUNDS FOR ELICITING OR ENHANCING IMMUNE REACTIVITY TO HER-2/*neu* PROTEIN FOR PREVENTION OR TREATMENT OF MALIGNANCIES IN WHICH THE HER-2/*neu* ONCOGENE IS ASSOCIATED

(iii) NUMBER OF SEQUENCES: 4

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: SEED and BERRY LLP
- (B) STREET: 6300 Columbia Center, 701 Fifth Avenue
- (C) CITY: Seattle
- (D) STATE: Washington
- (E) COUNTRY: USA
- (F) ZIP: 98104-7092

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE: 1-APR-1996
- (C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Sharkey, Richard G.
- (B) REGISTRATION NUMBER: 32,629
- (C) REFERENCE/DOCKET NUMBER: 920010.448C7

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (206) 622-4900
- (B) TELEFAX: (206) 682-6031

(2) INFORMATION FOR SEQ ID NO:1:

09167516.100698

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3768 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..3765

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GAG CTG GCG GCC TTG TGC CGC TGG GGG CTC CTC CTC GCC CTC TTG	48
Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu	
1 5 10 15	
CCC CCC GGA GCC GCG AGC ACC CAA GTG TGC ACC GGC ACA GAC ATG AAG	96
Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys	
20 25 30	
CTG CGG CTC CCT GCC AGT CCC GAG ACC CAC CTG GAC ATG CTC CGC CAC	144
Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His	
35 40 45	
CTC TAC CAG GGC TGC CAG GTG GTG CAG GGA AAC CTG GAA CTC ACC TAC	192
Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr	
50 55 60	
CTG CCC ACC AAT GCC AGC CTG TCC TTC CTG CAG GAT ATC CAG GAG GTG	240
Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val	
65 70 75 80	
CAG GGC TAC GTG CTC ATC GCT CAC AAC CAA GTG AGG CAG GTC CCA CTG	288
Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu	
85 90 95	
CAG AGG CTG CGG ATT GTG CGA GGC ACC CAG CTC TTT GAG GAC AAC TAT	336
Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr	
100 105 110	
GCC CTG GCC GTG CTA GAC AAT GGA GAC CCG CTG AAC AAT ACC ACC CCT	384
Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro	

09167516.100699

115	120	125	
GTC ACA GGG GCC TCC CCA GGA GGC CTG CGG GAG CTG CAG CTT CGA AGC Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser 130 135 140			432
CTC ACA GAG ATC TTG AAA GGA GGG GTC TTG ATC CAG CGG AAC CCC CAG Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln 145 150 155 160			480
CTC TGC TAC CAG GAC ACG ATT TTG TGG AAG GAC ATC TTC CAC AAG AAC Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn 165 170 175			528
AAC CAG CTG GCT CTC ACA CTG ATA GAC ACC AAC CGC TCT CGG GCC TGC Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys 180 185 190			576
CAC CCC TGT TCT CCG ATG TGT AAG GGC TCC CGC TGC TGG GGA GAG AGT His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser 195 200 205			624
TCT GAG GAT TGT CAG AGC CTG ACG CGC ACT GTC TGT GCC GGT GGC TGT Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys 210 215 220			672
GCC CGC TGC AAG GGG CCA CTG CCC ACT GAC TGC TGC CAT GAG CAG TGT Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys 225 230 235 240			720
GCT GCC GGC TGC ACG GGC CCC AAG CAC TCT GAC TGC CTG GCC TGC CTC Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu 245 250 255			768
CAC TTC AAC CAC AGT GGC ATC TGT GAG CTG CAC TGC CCA GCC CTG GTC His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val 260 265 270			816
ACC TAC AAC ACA GAC ACG TTT GAG TCC ATG CCC AAT CCC GAG GGC CGG Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg 275 280 285			864
TAT ACA TTC GGC GCC AGC TGT GTG ACT GCC TGT CCC TAC AAC TAC CTT Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu 290 295 300			912

09167516.100608

TCT ACG GAC GTG GGA TCC TGC ACC CTC GTC TGC CCC CTG CAC AAC CAA Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln 305 310 315 320	960
GAG GTG ACA GCA GAG GAT GGA ACA CAG CGG TGT GAG AAG TGC AGC AAG Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys 325 330 335	1008
CCC TGT GCC CGA GTG TGC TAT GGT CTG GGC ATG GAG CAC TTG CGA GAG Pro Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu 340 345 350	1056
GTG AGG GCA GTT ACC AGT GCC AAT ATC CAG GAG TTT GCT GGC TGC AAG Val Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys 355 360 365	1104
AAG ATC TTT GGG AGC CTG GCA TTT CTG CCG GAG AGC TTT GAT GGG GAC Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp 370 375 380	1152
CCA GCC TCC AAC ACT GCC CCG CTC CAG CCA GAG CAG CTC CAA GTG TTT Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Val Phe 385 390 395 400	1200
GAG ACT CTG GAA GAG ATC ACA GGT TAC CTA TAC ATC TCA GCA TGG CCG Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro 405 410 415	1248
GAC AGC CTG CCT GAC CTC AGC GTC TTC CAG AAC CTG CAA GTA ATC CGG Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg 420 425 430	1296
GGA CGA ATT CTG CAC AAT GGC GCC TAC TCG CTG ACC CTG CAA GGG CTG Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu 435 440 445	1344
GGC ATC AGC TGG CTG GGG CTG CGC TCA CTG AGG GAA CTG GGC AGT GGA Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly 450 455 460	1392
CTG GCC CTC ATC CAC CAT AAC ACC CAC CTC TGC TTC GTG CAC ACG GTG Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val 465 470 475 480	1440

09167516-100698

CCC TGG GAC CAG CTC TTT CGG AAC CCG CAC CAA GCT CTG CTC CAC ACT	1488
Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr	
485 490	
GCC AAC CGG CCA GAG GAC GAG TGT GTG GGC GAG GGC CTG GCC TGC CAC	1536
Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His	
500 505	
CAG CTG TGC GCC CGA GGG CAC TGC TGG GGT CCA GGG CCC ACC CAG TGT	1584
Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys	
515 520 525	
GTC AAC TGC AGC CAG TTC CTT CGG GGC CAG GAG TGC GTG GAG GAA TGC	1632
Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys	
530 535 540	
CGA GTA CTG CAG GGG CTC CCC AGG GAG TAT GTG AAT GCC AGG CAC TGT	1680
Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys	
545 550 555	
TTG CCG TGC CAC CCT GAG TGT CAG CCC CAG AAT GGC TCA GTG ACC TGT	1728
Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys	
565 570 575	
TTT GGA CCG GAG GCT GAC CAG TGT GTG GCC TGT GCC CAC TAT AAG GAC	1776
Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp	
580 585 590	
CCT CCC TTC TGC GTG GCC CGC TGC CCC AGC GGT GTG AAA CCT GAC CTC	1824
Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu	
595 600 605	
TCC TAC ATG CCC ATC TGG AAG TTT CCA GAT GAG GAG GGC GCA TGC CAG	1872
Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln	
610 615 620	
CCT TGC CCC ATC AAC TGC ACC CAC TCC TGT GTG GAC CTG GAT GAC AAG	1920
Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Lys	
625 630 635 640	
GGC TGC CCC GCC GAG CAG AGA GCC AGC CCT CTG ACG TCC ATC ATC TCT	1968
Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Ile Ser	
645 650 655	
GCG GTG GTT GGC ATT CTG CTG GTC GTG GTC TTG GGG GTG GTC TTT GGG	2016

Ala	Val	Val	Gly	Ile	Leu	Leu	Val	Val	Val	Leu	Gly	Val	Val	Phe	Gly	
			660						665						670	
ATC	CTC	ATC	AAG	CGA	CGG	CAG	CAG	AAG	ATC	CGG	AAG	TAC	ACG	ATG	CGG	2064
Ile	Leu	Ile	Lys	Arg	Arg	Gln	Gln	Lys	Ile	Arg	Lys	Tyr	Thr	Met	Arg	
			675					680							685	
AGA	CTG	CTG	CAG	GAA	ACG	GAG	CTG	GTG	GAG	CCG	CTG	ACA	CCT	AGC	GGA	2112
Arg	Leu	Leu	Gln	Glu	Thr	Glu	Leu	Val	Glu	Pro	Leu	Thr	Pro	Ser	Gly	
			690					695							700	
GCG	ATG	CCC	AAC	CAG	GCG	CAG	ATG	CGG	ATC	CTG	AAA	GAG	ACG	GAG	CTG	2160
Ala	Met	Pro	Asn	Gln	Ala	Gln	Met	Arg	Ile	Leu	Lys	Glu	Thr	Glu	Leu	
								710							720	
AGG	AAG	GTG	AAG	GTG	CTT	GGA	TCT	GGC	GCT	TTT	GGC	ACA	GTC	TAC	AAG	2208
Arg	Lys	Val	Lys	Val	Leu	Gly	Ser	Gly	Ala	Phe	Gly	Thr	Val	Tyr	Lys	
								725							735	
GGC	ATC	TGG	ATC	CCT	GAT	GGG	GAG	AAT	GTG	AAA	ATT	CCA	GTG	GCC	ATC	2256
Gly	Ile	Trp	Ile	Pro	Asp	Gly	Glu	Asn	Val	Lys	Ile	Pro	Val	Ala	Ile	
								745							750	
AAA	GTG	TTG	AGG	GAA	AAC	ACA	TCC	CCC	AAA	GCC	AAC	AAA	GAA	ATC	TTA	2304
Lys	Val	Leu	Arg	Glu	Asn	Thr	Ser	Pro	Lys	Ala	Asn	Lys	Glu	Ile	Leu	
								760							765	
GAC	GAA	GCA	TAC	GTG	ATG	GCT	GGT	GTG	GGC	TCC	CCA	TAT	GTC	TCC	CGC	2352
Asp	Glu	Ala	Tyr	Val	Met	Ala	Gly	Val	Gly	Ser	Pro	Tyr	Val	Ser	Arg	
								775							780	
CTT	CTG	GGC	ATC	TGC	CTG	ACA	TCC	ACG	GTG	CAG	CTG	GTG	ACA	CAG	CTT	2400
Leu	Leu	Gly	Ile	Cys	Leu	Thr	Ser	Thr	Val	Gln	Leu	Val	Thr	Gln	Leu	
								790							800	
ATG	CCC	TAT	GGC	TGC	CTC	TTA	GAC	CAT	GTC	CGG	GAA	AAC	CGC	GGA	CGC	2448
Met	Pro	Tyr	Gly	Cys	Leu	Leu	Asp	His	Val	Arg	Glu	Asn	Arg	Gly	Arg	
								805							815	
CTG	GGC	TCC	CAG	GAC	CTG	CTG	AAC	TGG	TGT	ATG	CAG	ATT	GCC	AAG	GGG	2496
Leu	Gly	Ser	Gln	Asp	Leu	Leu	Asn	Trp	Cys	Met	Gln	Ile	Ala	Lys	Gly	
								820							830	
ATG	AGC	TAC	CTG	GAG	GAT	GTG	CGG	CTC	GTA	CAC	AGG	GAC	TTG	GCC	GCT	2544
Met	Ser	Tyr	Leu	Glu	Asp	Val	Arg	Leu	Val	His	Arg	Asp	Leu	Ala	Ala	

00167516.100698

835										840										845											
CGG AAC GTG CTG GTC AAG AGT CCC AAC CAT GTC AAA ATT ACA GAC TTC																															2592
Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe																															
850																															
855																															
860																															
GGG CTG GCT CGG CTG CTG GAC ATT GAC GAG ACA GAG TAC CAT GCA GAT																															2640
Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp																															
865																															
870																															
875																															
880																															
GGG GGC AAG GTG CCC ATC AAG TGG ATG GCG CTG GAG TCC ATT CTC CGC																															2688
Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg																															
885																															
890																															
895																															
CGG CGG TTC ACC CAC CAG AGT GAT GTG TGG AGT TAT GGT GTG ACT GTG																															2736
Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val																															
900																															
905																															
910																															
TGG GAG CTG ATG ACT TTT GGG GCC AAA CCT TAC GAT GGG ATC CCA GCC																															2784
Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala																															
915																															
920																															
925																															
CGG GAG ATC CCT GAC CTG CTG GAA AAG GGG GAG CGG CTG CCC CAG CCC																															2832
Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro																															
930																															
935																															
940																															
945																															
950																															
955																															
960																															
965																															
970																															
975																															
980																															
985																															
990																															
995																															
1000																															
1005																															
1010																															
1015																															
1020																															
1025																															
1030																															
1035																															
1040																															
1045																															
1050																															
1055																															
1060																															
1065																															
1070																															
1075																															
1080																															
1085																															
1090																															
1095																															
1100																															
1105																															
1110																															
1115																															
1120																															
1125																															
1130																															
1135																															
1140																															
1145																															
1150																															
1155																															
1160																															
1165																															
1170																															
1175																															
1180																															
1185																															
1190																															
1195																															
1200																															
1205																															
1210																															
1215																															
1220																															
1225																															
1230																															
1235																															
1240																															
1245																															
1250																															
1255																															
1260																															
1265																															
1270																															
1275																															
1280																															
1285																															
1290																															
1295																															
1300																															
1305																															
1310																															
1315																															
1320																															
1325																															
1330																															
1335																															
1340																															
1345																															
1350																															
1355																															
1360																															
1365																															
1370																															
1375																															
1380																															
1385																															
1390																															
1395																															
1400																															
1405																															
1410																															
1415																															
1420																															
1425																															
1430																															
1435																															
1440																															
1445																															
1450																															
1455																															
1460																															
1465																															
1470																															
1475																															
1480																															
1485																															
1490																															
1495																															
1500																															
1505																															
1510																															
1515																															
1520																															
1525																															
1530																															
1535																															
1540																															
1545																															
1550																															
1555																															
1560																															
1565																															
1570																															
1575																															
1580																															
1585																															
1590																															
1595																															
1600																															
1605																															
1610																															
1615																															
1620																															
1625																															
1630																															
1635																															
1640																															
1645																															
1650																															
1655																															
1660																															
1665																															
1670																															
1675																															
1680																															
1685																															
1690																															
1695																															
1700																															
1705																															
1710																															
1715																															
1720																															
1725																															
1730																															
1735																															
1740																															
1745																															
1750																															
1755																															
1760																															
1765																															
1770																															
1775																															
1780																															
1785																															
1790																															
1795																															
1800																															
1805																															
1810																															
1815																															
1820																															
1825																															
1830																															
1835																															
1840																															
1845																															
1850																															
1855																															
1860																															
1865																															
1870																															
1875																															
1880																															
1885																															
1890																															
1895																															
1900																															
1905																															
1910																															
1915																															
1920																															
1925																															
1930																															
1935																															
1940																															
1945																															
1950																															
1955																															
1960																															
1965																															
1970																															
1975																															
1980																															
1985																															
1990																															
1995																															
2000																															
2005																															
2010																															
2015																															
2020																															
2025																															
2030																															
2035																															
2040																															
2045																															
2050																															
2055																															
2060																															
2065																															
2070																															
2075																															
2080																															
2085																															
2090																															
2095																															
2100																															
2105																															
2110																															
2115																															
2120																															
2125																															
2130																															
2135																															
2140																															
2145																															
2150																															
2155																															
2160																															
2165																															
2170																															
2175																															
2180																															
2185																															
2190																															
2195																															
2200																															
2205																															
2210																															
2215																															
2220																															
2225																															
2230																															
2235																															
2240																															
2245																															
2250																															
2255																															
2260																															</

GTA CCC CAG CAG GGC TTC TTC TGT CCA GAC CCT GCC CCG GGC GCT GGG Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly 1025 1030 1035 1040	3120
GGC ATG GTC CAC CAC AGG CAC CGC AGC TCA TCT ACC AGG AGT GGC GGT Gly Met Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly 1045 1050 1055	3168
GGG GAC CTG ACA CTA GGG CTG GAG CCC TCT GAA GAG GAG GCC CCC AGG Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Ala Pro Arg 1060 1065 1070	3216
TCT CCA CTG GCA CCC TCC GAA GGG GCT GGC TCC GAT GTA TTT GAT GGT Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly 1075 1080 1085	3264
GAC CTG GGA ATG GGG GCA GCC AAG GGG CTG CAA AGC CTC CCC ACA CAT Asp Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His 1090 1095 1100	3312
GAC CCC AGC CCT CTA CAG CGG TAC AGT GAG GAC CCC ACA GTA CCC CTG Asp Pro Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu 1105 1110 1115 1120	3360
CCC TCT GAG ACT GAT GGC TAC GTT GCC CCC CTG ACC TGC AGC CCC CAG Pro Ser Glu Thr Asp Gly Tyr Val Ala Pro Leu Thr Cys Ser Pro Gln 1125 1130 1135	3408
CCT GAA TAT GTG AAC CAG CCA GAT GTT CGG CCC CAG CCC CCT TCG CCC Pro Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro 1140 1145 1150	3456
CGA GAG GGC CCT CTG CCT GCT GCC CGA CCT GCT GGT GCC ACT CTG GAA Arg Glu Gly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu 1155 1160 1165	3504
AGG CCC AAG ACT CTC TCC CCA GGG AAG AAT GGG GTC GTC AAA GAC GTT Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val 1170 1175 1180	3552
TTT GCC TTT GGG GGT GCC GTG GAG AAC CCC GAG TAC TTG ACA CCC CAG Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln 1185 1190 1195 1200	3600

GGA GGA GCT GCC CCT CAG CCC CAC CCT CCT CCT GCC TTC AGC CCA GCC 3648
 Gly Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala
 1205 1210 1215

TTC GAC AAC CTC TAT TAC TGG GAC CAG GAC CCA CCA GAG CGG GGG GCT 3696
 Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala
 1220 1225 1230

CCA CCC AGC ACC TTC AAA GGG ACA CCT ACG GCA GAG AAC CCA GAG TAC 3744
 Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr
 1235 1240 1245

CTG GGT CTG GAC GTG CCA GTG TGA 3768
 Leu Gly Leu Asp Val Pro Val
 1250 1255

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1255 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu
 1 5 10 15

Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys
 20 25 30

Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His
 35 40 45

Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr
 50 55 60

Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val
 65 70 75 80

Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu
 85 90 95

09167516-100698

Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr
 100 105 110
 Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro
 115 120 125
 Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser
 130 135 140
 Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln
 145 150 155 160
 Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn
 165 170 175
 Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys
 180 185 190
 His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser
 195 200 205
 Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys
 210 215 220
 Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys
 225 230 235 240
 Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu
 245 250 255
 His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val
 260 265 270
 Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg
 275 280 285
 Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu
 290 295 300
 Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln
 305 310 315 320
 Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys
 325 330 335

09167516-100698

Pro Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu
340 345 350

Val Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys
355 360 365

Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp
370 375 380

Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe
385 390 395 400

Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro
405 410 415

Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg
420 425 430

Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu
435 440 445

Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly
450 455 460

Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val
465 470 475 480

Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr
485 490 495

Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His
500 505 510

Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys
515 520 525

Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys
530 535 540

Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys
545 550 555 560

Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys
565 570 575

09167516.100000

Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp
580 585 590

Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu
595 600 605

Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln
610 615 620

Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys
625 630 635 640

Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Ile Ser
645 650 655

Ala Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly
660 665 670

Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg
675 680 685

Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly
690 695 700

Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu
705 710 715 720

Arg Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys
725 730 735

Gly Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile
740 745 750

Lys Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu
755 760 765

Asp Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg
770 775 780

Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu
785 790 795 800

Met Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg
805 810 815

09167516-100698

Leu Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly
820 825 830

Met Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala
835 840 845

Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe
850 855 860

Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp
865 870 875 880

Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg
885 890 895

Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val
900 905 910

Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala
915 920 925

Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro
930 935 940

Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met
945 950 955 960

Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe
965 970 975

Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu
980 985 990

Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu
995 1000

Leu Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr Leu
1010 1015 1020

Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly
1025 1030 1035 1040

Gly Met Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly
1045 1050 1055

09167516-100698

Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Ala Pro Arg
1060 1065 1070

Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly
1075 1080 1085

Asp Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His
1090 1095 1100

Asp Pro Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu
1105 1110 1115 1120

Pro Ser Glu Thr Asp Gly Tyr Val Ala Pro Leu Thr Cys Ser Pro Gln
1125 1130 1135

Pro Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro
1140 1145 1150

Arg Glu Gly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu
1155 1160 1165

Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val
1170 1175 1180

Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln
1185 1190 1195 1200

Gly Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala
1205 1210 1215

Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala
1220 1225 1230

Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr
1235 1240 1245

Leu Gly Leu Asp Val Pro Val
1250 1255

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 48 base pairs

(B) TYPE: nucleic acid

09167516.100698

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TCTGGCGCGC TGGATGACGA TGACAAGAAA CGACGGCAGC AGAAGATC

48

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TGAATTCTCG AGTCATTACA CTGGCAGTC CAGACCCAG

39

09167516.100698

Claims

1. A polypeptide encoded by a DNA sequence selected from:

(a) nucleotides 2026 through 3765 of SEQ ID NO:1; and

(b) DNA sequences that hybridize to a nucleotide sequence complementary to nucleotides 2026 through 3765 of SEQ ID NO:1 under moderately stringent conditions, wherein the DNA sequence encodes a polypeptide that produces an immune response to HER-2/*neu* protein.

2. A polypeptide having the amino acid sequence of SEQ ID NO:2 from lysine, amino acid 676, through valine, amino acid 1255, or a variant thereof that produces at least an equivalent immune response.

3. A polypeptide according to claim 2 having the amino acid sequence of SEQ ID NO:2 from amino acid 676 through amino acid 1255.

4. A composition comprising a polypeptide according to any one of claims 1, 2 or 3, in combination with a pharmaceutically acceptable carrier or diluent.

5. A nucleic acid molecule directing the expression of a polypeptide according to any one of claims 1, 2 or 3.

6. A viral vector directing the expression of a polypeptide according to any one of claims 1, 2 or 3.

7. A method for eliciting or enhancing an immune response to HER-2/*neu* protein, comprising administering to a

09167516-100668

warm-blooded animal in an amount effective to elicit or enhance said response a polypeptide according to any one of claims 1, 2 or 3, or a nucleic acid molecule according to claim 5, or a viral vector according to claim 6.

8. A method according to claim 7 wherein the step of administering comprises transfecting cells of the animal *ex vivo* with the nucleic acid molecule and subsequently delivering the transfected cells to the animal.


9. A method according to claim 7 wherein the step of administering comprises infecting cells of the animal *ex vivo* with the viral vector and subsequently delivering the infected cells to the animal.


09167516.100000


COMPOUNDS FOR ELICITING OR ENHANCING IMMUNE REACTIVITY
TO HER-2/*neu* PROTEIN FOR PREVENTION OR TREATMENT OF
MALIGNANCIES IN WHICH THE HER-2/*neu* ONCOGENE IS ASSOCIATED


Abstract of the Disclosure

Compounds and compositions for eliciting or enhancing immune reactivity to HER-2/*neu* protein are disclosed. The compounds include polypeptides and nucleic acid molecules encoding such peptides. The compounds may be used for the prevention or treatment of malignancies in which the HER-2/*neu* oncogene is associated.

**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

DECLARATION AND POWER OF ATTORNEY

As the below-named inventors, we declare that:

Our residences, post office addresses, and citizenships are as stated below under our names.

We believe we are the original, first and joint inventors of the invention entitled "COMPOUNDS FOR ELICITING OR ENHANCING IMMUNE REACTIVITY TO HER-2/*neu* PROTEIN FOR PREVENTION OR TREATMENT OF MALIGNANCIES IN WHICH THE HER-2/*neu* ONCOGENE IS ASSOCIATED," which is described and claimed in the specification and claims of C-I-P Application No. 08/625,101, which we filed in the United States Patent and Trademark Office on April 1, 1996, and for which a patent is sought; and that this application in part discloses and claims subject matter disclosed in our earlier-filed Application No. 08/414,417, entitled "IMMUNE REACTIVITY TO HER-2/*neu* PROTEIN FOR DIAGNOSIS AND TREATMENT OF MALIGNANCIES IN WHICH THE HER-2/*neu* ONCOGENE IS ASSOCIATED," which we filed in the United States Patent and Trademark Office on March 31, 1995.

We have reviewed and understand the contents of the above-entitled specification, including the claims, as amended by any amendment specifically referred to herein (if any).

We acknowledge our duty to disclose information of which we are aware which is material to the examination of this application in accordance with 37 C.F.R. § 1.56(a), including material information which occurred between the filing date of said earlier-filed pending application and the filing date of the C-I-P application.

We hereby appoint RICHARD W. SEED, Registration No. 16,557; ROBERT J. BAYNHAM, Registration No. 22,846; EDWARD W. BULCHIS, Registration No. 26,847; GEORGE C. RONDEAU, JR., Registration No. 28,893; DAVID H. DEITS, Registration No. 28,066; WILLIAM O. FERRON, JR., Registration No. 30,633; PAUL T. MEIKLEJOHN, Registration No. 26,569; DAVID J. MAKI, Registration No. 31,392; RICHARD G. SHARKEY, Registration No. 32,629; DAVID V. CARLSON, Registration No. 31,153; MAURICE J. PIRIO, Registration No. 33,273; KARL R. HERMANN, Registration No. 33,507; DAVID D. McMASTERS, Registration No. 33,963; ROBERT IANNUCCI, Registration No. 33,514; JOSHUA KING, Registration No. 35,570; MICHAEL J. DONOHUE, Registration No. 35,859; LORRAINE LINFORD, Registration No. 35,939; KEVIN J. CANNING, Registration No. 35,470; CHRISTOPHER J. DALEY-WATSON, Registration No. 34,807; STEVEN D. LAWRENZ, Registration No. 37,376; ROBERT G. WOOLSTON, Registration No. 37,263; CLARENCE T. TEGREENE, Registration No. 37,951; ELLEN M. BIERMAN, Registration No. 38,079; BRYAN A. SANTARELLI, Registration No. 37,560; MICHAEL L. KIKLIS, Registration No. 38,939;

09167516-100698


CAROL NOTTENBURG, Registration No. 39,317; CRAIG S. JEPSON, Registration No. 33,517; PAUL T. PARKER, Registration No. 38,264; JOHN C. STEWART, Registration No. 40,188; ROBERT W. BERGSTROM, Registration No. 39,906; HARRY K. AHN, Registration No. 40,243; and DAVID W. PARKER, Registration No. 37,414, comprising the firm of SEED and BERRY LLP, 6300 Columbia Center, Seattle, Washington 98104-7092, as our attorneys to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. Please direct all telephone calls to Richard G. Sharkey at (206) 622-4900 and telecopies to (206) 682-6031.


We further declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that the making of willfully false statements and the like is punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and may jeopardize the validity of any patent issuing from this patent application.


 Martin A. Cheever

Date 6/20/96
 Residence : City of Mercer Island, County of King
 State of Washington
 Citizenship : United States of America
 P.O. Address : 6825 - 83rd Avenue S.E.
 Mercer Island, Washington 98040


SUBSCRIBED AND SWORN TO BEFORE ME
 this 20 day of June, 19 96


 Jim W. Ponder comm exp - 3/1/2000
 Notary Public in and for the State
 of Washington, King County, Seattle


 Mary L. Disis

Date 6/28/96
 Residence : City of Renton, County of King
 State of Washington
 Citizenship : United States of America
 P.O. Address : 17850 S.E. 192nd Drive
 Renton, Washington 98055

SUBSCRIBED AND SWORN TO BEFORE ME
 this 28 day of June, 19 96


 Jim W. Ponder comm exp - 3/1/2000
 Notary Public in and for the State
 of Washington, King County, Seattle